

Summary (Part 1 Pure)

Arithmetic

Definitions;

Sum = One number plus another

Difference = One number minus another

Product = One number times another

Quotient = One number divided by another

A number is the Product of its Factors

Primes are Numbers with no Factors except 1 and itself

HCF (Highest Common Factor) = Highest Factor that is Common to all numbers of a group

LCM (Lowest Common Multiplier) = Lowest number that has all numbers of a group as Factors

Numerator is the Number at top of Fraction

Denominator is the Number at bottom of Fraction

Reciprocal = 1 Divided by the Number

Factorial is the Product of all Numbers from 1 to the Number and is written with ! For example $4! = 1 \times 2 \times 3 \times 4$

Ratio is the Comparison of 2 or more Numbers. For example $15 : 5$ has the same Ratio as $3 : 1$

Square of a Number = Number times itself, written as N^2 . For example $5^2 = 25$

Square Root of a Number times itself = The Number. Square Root is written as \sqrt{N} . For example $\sqrt{25} = \pm 5$

Index, or Power = Number of times a Number is multiplied by itself. For example $5^3 = 5 \times 5 \times 5$ has the index of 3

Scientific Notation = Number expressed as a number between 0 and 10 times powers of 10

Binary = Number expressed in 2 digits (0 & 1)

Octal = Number expressed in 8 digits (0 - 7)

Hexadecimal = Number expressed in 16 digits (0 - 9 and A - F)

Hex(abcd) = Decimal $(d + c \times 16 + b \times 16^2 + a \times 16^3)$
 $2^{10} = 1024$

Algebra

Definitions;

Irrational functions are Functions containing a square root, or cube root etc

Equations are statements that two functions are equal

Simultaneous equations are a set of Equations connecting two or more unknowns

Greek letter sigma Σ means "Sum of Terms Like"

Coefficient. For example in the function $7x^2 - 5x + 3$, the coefficient of x^2 is 7 and the coefficient of x is -5

$(-a) \text{ times } (-b) = + ab$

$a^m \text{ times } a^n = a^{m+n}$

$(a^m)^n = a^{mn}$

$a^0 = 1$ and $a^1 = a$ and $a^{-n} = 1/a^n$ and $a^{(1/n)} = \sqrt[n]{a}$

$(a + b)(c + d) = a(c + d) + b(c + d) = ac + ad + bc + bd$

To factorize $ax^2 + bx + c$

If ac (ie a times c) is -ive, look for factors of ac whose sum = b

If ac is +ive, look for factors whose difference = $\pm b$

$x^2 - a^2 = (x + a)(x - a)$

$(x \pm 1)$ is a factor of $x^3 \pm 1$

$a_0 + a_1x + a_2x^2 + \dots + a_nx^n$ can be divided by $b_0 + b_1x + b_2x^2 + \dots + b_{n-r}x^{n-r}$ to get **Quotient** and **Remainder**

Method is similar to Arithmetical Long Division. Include missing terms using zero as the coefficient

For example $(ax^2 + bx + c)/(x + d) = [ax + (bd - ad)] + [(c - bd + ad^2)/(x + d)]$

Divide $F(x)$ by $(x - a)$ and the **Remainder** is $F(a)$

$F(x) / [(x + \alpha_1)(x + \alpha_2)(x + \alpha_3)] = A_1/(x + \alpha_1) + A_2/(x + \alpha_2) + A_3/(x + \alpha_3)$ where $A_1 = F(-\alpha_1) / [(\alpha_2 - \alpha_1)(\alpha_3 - \alpha_1)]$ etc

If the Numerator is same or higher power than the Denominator, divide it by the Denominator first.

In each fraction, the Numerator contains x to one power less than the Denominator

Two equal factors $F(x) / [(x + \alpha_1)^2(x + \alpha_3)] = A_1/(x + \alpha_1) + A_2/(x + \alpha_1)^2 + A_3/(x + \alpha_3)$

$1/(a + \sqrt{b}) = (a - \sqrt{b})/(a^2 - b)$ and $1/(a - \sqrt{b}) = (a + \sqrt{b})/(a^2 - b)$ this puts the irrational term in the numerator

$i^2 = -1$

$(a + ib)(a - ib) = a^2 + b^2$ and $1/(a + ib) = (a - ib)/(a^2 + b^2)$ and $1/(a - ib) = (a + ib)/(a^2 + b^2)$. Puts in form $c + id$

Solution to the quadratic $Ax^2 + Bx + C = 0$ is $x = [-B \pm \sqrt{B^2 - 4AC}] / 2A$

$\alpha_1 + \alpha_2 = -B/A$ and $\alpha_1 \alpha_2 = C/A$ where α_1 and α_2 are the two solutions

If $4AC > B^2$ then the two solutions are a conjugate pair, $\alpha + i\beta$ and $\alpha - i\beta$

An equation has as many solutions as the highest power of x after rationalizing. A quadratic has 2, a cubic has 3.

Geometry

Definitions;

Angles, one revolution is 360 degrees = 2π radians, a Right Angle is 90 degrees = $\pi/2$ radians

Equilateral Triangle has all sides equal

Isosceles Triangle has two angles equal

Similar Triangles are Two Triangles with same angles. The Ratio between the sides is the same in both.

Congruent Triangles are Two Triangles exactly the same

“Normal to”, “Orthogonal to” and “Perpendicular to” mean “at right angles to”

Hypotenuse is the Side of a Triangle opposite a right angle

Tangent is a line that just touches a curve and is parallel to the curve at that point.

Length of arc = radius times angle in radians $s = r\theta$

Conditions for congruent triangles either (i) 3 sides same, or (ii) 2 sides and the angle between them the same, or (iii) 2 angles and a corresponding side the same or (iv) hypotenuse and one other side the same.

Triangle **Sum of angles = 180°**

Area = $(\frac{1}{2})$ Base x Height

Pythagoras (for a Right Angled Triangle) **$a^2 + b^2 = c^2$**

Examples $3^2 + 4^2 = 5^2$, $6^2 + 8^2 = 10^2$, $12^2 + 5^2 = 13^2$


Medians meet at a point, so do Angle bisectors, so do lines from each apex perpendicular to opposite side, so do perpendiculars from mid points of sides

Circle **Circumference = $\pi D = 2\pi R$** and **Area = πR^2** where D is the Diameter and R the Radius

Rectangle or Parallelogram **Area = Base x Height** (Height is measured normal to base)

Trigonometry

$\sin \theta = a/c$ and **$\cos \theta = b/c$** and **$\tan \theta = a/b$**
 $\operatorname{cosec} \theta = 1/\sin \theta$ and **$\operatorname{sec} \theta = 1/\cos \theta$** and **$\cot \theta = 1/\tan \theta$**
 $\sin \theta / \cos \theta = \tan \theta$




$\sin^2 \theta + \cos^2 \theta = 1$ and **$\tan^2 \theta + 1 = \sec^2 \theta$**


$\sin(90^\circ - \theta) = \cos \theta$ and **$\tan(90^\circ - \theta) = \cot \theta$**

$\sin(-\theta) = -\sin \theta$ and **$\cos(-\theta) = +\cos \theta$** and **$\tan(-\theta) = -\tan \theta$**
 $\sin(180^\circ - \theta) = +\sin \theta$ and **$\cos(180^\circ - \theta) = -\cos \theta$** and **$\tan(180^\circ - \theta) = -\tan \theta$**
 $\sin(180^\circ + \theta) = -\sin \theta$ and **$\cos(180^\circ + \theta) = -\cos \theta$** and **$\tan(180^\circ + \theta) = +\tan \theta$**


CAST Angles (-90° to 0°) **Cos + ive, Sin and Tan - ive**
 Angles (0° to 90°) **All + ive**
 Angles (90° to 180°) **Sin + ive, Cos and Tan - ive**
 Angles (180° to 270°) **Tan + ive, Sin and Cos - ive**



If θ is small and in radians then **$\sin \theta = \theta$**
 and **$\tan \theta = \theta$** and **$\cos \theta = 1 - (\frac{1}{2})\theta^2$**



For other angles, see the diagrams



$$\begin{aligned} \sin(A + B) &= \sin A \cos B + \cos A \sin B \\ \cos(A + B) &= \cos A \cos B - \sin A \sin B \\ \tan(A + B) &= (\tan A + \tan B) / (1 - \tan A \tan B) \\ \sin(2A) &= 2 \sin A \cos A \\ \cos(2A) &= \cos^2 A - \sin^2 A \\ \tan(2A) &= 2 \tan A / (1 - \tan^2 A) \end{aligned}$$


$$\begin{aligned} \sin A \cos B &= (1/2) [\sin(A + B) + \sin(A - B)] \\ \cos A \cos B &= (1/2) [\cos(A + B) + \cos(A - B)] \\ \sin A \sin B &= (1/2) [\cos(A - B) - \cos(A + B)] \\ \sin A + \sin B &= 2 \sin[(1/2)(A + B)] \cos[(1/2)(A - B)] \\ \sin A - \sin B &= 2 \cos[(1/2)(A + B)] \sin[(1/2)(A - B)] \\ \cos A + \cos B &= 2 \cos[(1/2)(A + B)] \cos[(1/2)(A - B)] \\ \cos A - \cos B &= -2 \sin[(1/2)(A + B)] \sin[(1/2)(A - B)] \\ \sin^2 A - \sin^2 B &= \sin(A + B) \sin(A - B) \\ \cos^2 A - \cos^2 B &= -\sin(A + B) \sin(A - B) \\ \cos^2 A - \sin^2 B &= \cos(A + B) \cos(A - B) \end{aligned}$$

Triangle

$$\begin{aligned} \text{Area} &= (1/2) ab \sin C \quad \text{and} \quad \text{Area} = \sqrt{[s(s-a)(s-b)(s-c)]} \quad \text{where} \quad s = (1/2)(a + b + c) \\ a/\sin A &= b/\sin B = c/\sin C \quad \text{and} \quad c^2 = a^2 + b^2 - 2ab \cos C \end{aligned}$$

Co-ordinate Geometry

Cartesian co-ordinates, points are shown by x and y
 Polar co-ordinates, points are shown by r and θ
 $r = \sqrt{x^2 + y^2}$ and $\tan \theta = y/x$ and $x = r \cos \theta$ and $y = r \sin \theta$



$$\begin{aligned} \text{Straight line, slope } m & \quad y = mx + c \\ \text{Line through } (x_1, y_1) \text{ and } (x_2, y_2) & \quad y_1 = m x_1 + c \text{ and } y_2 = m x_2 + c \text{ Solve for } m \text{ and } c \\ \text{Angle between two lines } \tan \theta &= (m_1 - m_2) / (1 + m_1 m_2) \\ \text{2 lines cross orthogonally if } & \quad m_1 m_2 = -1 \\ \text{Circle, centre at origin} & \quad x^2 + y^2 = a^2 \\ \text{Circle, centre at } (g, h), \text{ radius } a & \quad (x - g)^2 + (y - h)^2 = a^2 \\ \text{Ellipse, centre at origin} & \quad x^2/a^2 + y^2/b^2 = 1 \\ \text{Parabola} & \quad y^2 = 4ax \\ \text{Hyperbola} & \quad xy = c^2 \text{ or } x^2/a^2 - y^2/b^2 = 1 \end{aligned}$$

Logarithms

$$\begin{aligned} \text{By definition; } \log_a m &= x \text{ where } a^x = m \\ \text{Hence } \log_a m + \log_a n &= \log_a(mn) \\ \log_a m - \log_a n &= \log_a(m/n) \\ n \log_a m &= \log_a m^n \\ \log_b m &= \log_a m / \log_a b \end{aligned}$$

Binominal

$$(x + a)^n = x^n + n a x^{n-1} + [n(n-1)/2!] a^2 x^{n-2} + [n(n-1)(n-2)/3!] a^3 x^{n-3} \dots + n! / [(n-r)! r!] a^r x^{n-r} + \dots + a^n$$

Matrices

Data can be displayed and manipulated in short hand in the form of Matrices.

Eg $a_1 x + a_2 y + a_3 z + a_4 = 0$ can be written (and solved) in Matrix form
 $b_1 x + b_2 y + b_3 z + b_4 = 0$
 $c_1 x + c_2 y + c_3 z + c_4 = 0$
 (add or subtract lines to get coefficients 2 and 3 = 0 to solve for x etc)

$$\begin{array}{cccc|c} a_1 & a_2 & a_3 & a_4 & |x| = 0 \\ b_1 & b_2 & b_3 & b_4 & |y| \\ c_1 & c_2 & c_3 & c_4 & |z| \\ & & & & |1| \end{array}$$

Determinants

$$\begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1 b_2 - a_2 b_1$$

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = a_1 b_2 c_3 - a_1 b_3 c_2 - a_2 b_1 c_3 + a_2 b_3 c_1 + a_3 b_1 c_2 - a_3 b_2 c_1$$

terms in sequence abc and numbers in sequence 123123 positive, others negative

Series

Definitions;

Arithmetical Progression AP is a Series with the same difference between all adjacent terms

Geometrical Progression GP is a Series with the same ratio between all adjacent terms

Sum S of AP, 1st term a, difference d, n terms. **Add first term to last term**, 2nd term to 2nd last etc
 hence $2S = n [\{a\} + \{a + (n - 1) d\}]$

Sum S of GP, 1st term a, ratio of terms p, n terms.

Then **Series - p times Series = first term + last term** hence $S = a (1 - p^n)/(1 - p)$

Sum of first n numbers is an AP = $n(n + 1)/2$

Sum of first n squares = $(1/6)n(n + 1)(2n + 1)$

Sum of first n cubes = $[(n + 1) n/2]^2$

Calculus

Definitions;

Differential of y, written dy/dx is Slope at x

Integral of y, written $\int y dx$ is the sum of areas of height y and width dx

$$d/dx[ax^n] = a n x^{n-1} \quad \text{and} \quad \int a x^n dx = a x^{n+1}/(n+1) + c$$

Integration between limits. The value of the Integral with the final value of x minus its value with the initial x

In polar co-ordinates, $dy/dx = (\sin \theta dr/d\theta + r \cos \theta)/(\cos \theta dr/d\theta - r \sin \theta)$

In polar co-ordinates, Sum of areas = $\int (\frac{1}{2}) r^2 d\theta$

Differential of a sum

$$d/dx (u + v) = du/dx + dv/dx$$

Differential of a product

$$d/dx (u v) = v du/dx + u dv/dx$$

Differential of a Fraction

$$d/dx (u/v) = \{ v du/dx - u dv/dx \} / v^2$$

Differential with a change of variable

$$dy/dx = dy/du \cdot du/dx$$

ln is natural logarithm (ie Log to base e) where $e = 1 + 1/1! + 1/2! + 1/3! + \dots$ to infinity

By definition, if $\ln(m) = x$ then $e^x = m$

$$d/dx (e^x) = e^x \quad \text{and} \quad \int (1/x) dx = \ln(x) + c \quad \text{and} \quad d/dx [\ln(x)] = 1/x$$

$$a^x = e^{x \ln(a)}$$

$$d/dx (\sin x) = \cos x \quad \text{and} \quad d/dx (\cos x) = -\sin x \quad \text{and} \quad d/dx (\tan x) = \sec^2 x$$

$$d/dx \{ \text{Arc Sin } (x/a) \} = 1 / \sqrt{a^2 - x^2} \quad \text{and} \quad d/dx \{ \text{Arc Cos } (x/a) \} = -1 / \sqrt{a^2 - x^2}$$

$$d/dx \{ \text{Arc Tan } (x/a) \} = a / (a^2 + x^2)$$

MacLaurim's Theorem

Let $f(x) = a_0 + a_1x/1! + a_2x^2/2! + \dots + a_r x^r/r! +$

Write $f_r(0)$ to mean r th differential of $f(x)$ with x then made zero, hence $a_r = f_r(0)$

$$f(x) = f(0) + f_1(0) x/1! + f_2(0) x^2/2! + f_3(0) x^3/3! + \dots + f_r(0) x^r/r! +$$

The series for many functions can be written down, eg

$$\sin x = x - x^3/3! + x^5/5! - x^7/7! \dots$$

$$\cos x = 1 - x^2/2! + x^4/4! - x^6/6! \dots$$

$$e^x = 1 + x/1! + x^2/2! + x^3/3! + x^4/4! \dots$$

$$\ln(1+x) = x - x^2/2 + x^3/3 - x^4/4 \dots$$

Taylor's Theorem

$$f(x) = f(a) + (x-a)f_1(a) + \dots + [(x-a)^r/r!] f_r(a) + \dots$$

Hyperbolic Functions

Expand $\cos n\theta + i \sin n\theta$ by MacLaurim's Theorem and the result is the expansion of $e^{in\theta}$

$$\cos n\theta + i \sin n\theta = e^{in\theta} = [\cos \theta + i \sin \theta]^n$$

$$\cos \theta + i \sin \theta = e^{i\theta} \quad \text{and} \quad \cos \theta - i \sin \theta = e^{-i\theta}$$

$$\cos \theta = \{e^{i\theta} + e^{-i\theta}\} / 2 \quad \text{and} \quad \sin \theta = \{e^{i\theta} - e^{-i\theta}\} / 2i$$

By definition, Cosh and Sinh are these values of Cos and Sin without the i

$$\cosh \theta = \{e^\theta + e^{-\theta}\} / 2 \quad \text{and} \quad \sinh \theta = \{e^\theta - e^{-\theta}\} / 2$$

$$\tanh \theta = (\sinh \theta) / (\cosh \theta) \quad \text{and} \quad \operatorname{sech} \theta = 1/\cosh \theta$$

$$\operatorname{cosech} \theta = 1/\sinh \theta \quad \text{and} \quad \coth \theta = 1/\tanh \theta$$

$$\cosh^2 \theta - \sinh^2 \theta = 1 \quad \text{and} \quad 1 - \tanh^2 \theta = \operatorname{sech}^2 \theta$$

$$\sinh 2x = 2 \sinh x \cosh x \quad \text{and} \quad \cosh 2x = \cosh^2 x + \sinh^2 x$$

$$d/dx (\sinh x) = \cosh x \quad \text{and} \quad d/dx (\cosh x) = \sinh x \quad \text{and} \quad d/dx (\tanh x) = \operatorname{sech}^2 x$$

Methods for Integration

In General Look for a substitution that will simplify the integral

$\int F(ax \pm b) dx$ indicates the substitution $u = (ax \pm b)$ thus $du = a dx$

eg $\int [\sin(x+a)] dx$ Put $u = x+a$ thus Integral = $\int [\sin(u)] du = -\cos(u) + \text{constant}$

$\int [1/(x^2 + a^2)] dx$ indicates the substitution $x = a \tan(u)$ and $dx = a \sec^2(u) du$ or $x = a \sinh(u)$ and $dx = a \cosh(u) du$

eg $\int [1/(x^2 + a^2)] dx$ Put $x = a \tan(u)$ this leads to $(1/a) \int du = u/a + \text{constant}$

Fractions If the denominator factorizes, Split into Partial Fractions;

$$\int [1/\{(x \pm a)(x \pm b)\}] dx = \int [A/(x \pm a)] dx + \int [B/(x \pm b)] dx$$

$$\text{eg } \int [1/(x^2 - a^2)] dx = \int [(1/2a)/(x-a)] dx - \int [(1/2a)/(x+a)] dx = (1/2a)[\ln(x-a) - \ln(x+a)] + \text{constant}$$

Integrals of Square Roots (Remember $1 - \sin^2 u = \cos^2 u$, $1 + \sinh^2 u = \cosh^2 u$, and $\cosh^2 u - 1 = \sinh^2 u$)

$\int [1/\sqrt{a^2 - x^2}] dx$ indicates the substitution $x = a \sin(u)$ thus $dx = a \cos(u) du$

$$\int [1/\sqrt{a^2 - x^2}] dx = \int [1/a \cos(u)] a \cos(u) du = \int du = u + \text{constant}$$

$\int [1/\sqrt{a^2 + x^2}] dx$ indicates the substitution $x = a \sinh(u)$ or $x = a \tan(u)$

$$\int [1/\sqrt{x^2 + a^2}] dx \text{ Put } x = a \sinh(u) \text{ this leads to } \int du = u + \text{constant}$$

$\int [1/\sqrt{x^2 - a^2}] dx$ indicates the substitution $x = a \cosh(u)$

$$\int [1/\sqrt{x^2 - a^2}] dx \text{ Put } x = a \cosh u \text{ leads to } \int du = u + \text{constant}$$

$\int [1/\sqrt{ax^2 + bx + c}] dx$ Remove the x term,

Put $a[(x+p)^2 + q] = ax^2 + bx + c$ Equate coefficients to solve for p and q , Put $u = x+p$ and $r^2 = q$.

This leads to $(1/\sqrt{a}) \int [1/\sqrt{u^2 \pm r^2}] du$ As above put $u = r \sinh v$ or $u = r \cosh v$

$\int \sqrt{(a+bx)/(c-dx)} dx$ where $a, b, c,$ and d are all +ve. Make coefficients of $x = 1$

$I = \sqrt{(b/d)} \int \sqrt{(a/b+x)/(c/d-x)} dx$. Put $x = p + q \sin \theta$ hence $dx = q \cos \theta d\theta$. Find values for p and q to get

$k \int \sqrt{[(1 + \sin \theta)/(1 - \sin \theta)] \cos \theta} d\theta$ Evaluate k . Multiply top and bottom by $(1 + \sin \theta)$ to get $k \int [1 + \sin \theta] d\theta$

Trigonometrical integrals

(i) Put in form $\int F(\cos x) \sin x \, dx$, or $\int F(\sin x) \cos x \, dx$ or $\int F(\tan x) \sec^2 x \, dx$ and Substitute to get $\int F(u) \, du$
 Similarly for hyperbolics Eg $\int \sinh^3 x \, dx = \int (\cosh^2 x - 1) \sinh x \, dx = \frac{1}{3} \cosh^3 x - \cosh x + \text{constant}$
 or (ii) Try the substitution $u = \tan(x)$ since $dx = du/(1 + u^2)$
 or (iii) Try substituting $t = \tan(x/2)$ since $dx = 2 \, dt/(1 + t^2)$, $\sin(x) = 2t/(1 + t^2)$ and $\cos(x) = (1 - t^2)/(1 + t^2)$
 $\int [1/(a \sin x + b \cos x + c)] dx$ indicates the substitution $t = \tan(x/2)$
 $\int \cos^2(x) \, dx$ and $\int \sin^2(x) \, dx$ indicate the substitution $u = 2x$ since $\cos^2(x) = \frac{1}{2} [\cos(u) + 1]$ and $dx = (\frac{1}{2}) \, du$

1/D Method

The operator D is defined as d/dx . Thus $D(y) = dy/dx$ hence $(D + a)(D + b)(y) = D^2(y) + (a + b)D(y) + ab y$

$$D^{-1}(y) = \int y \, dx$$

$$D^n (e^{ax} V) = e^{ax} (D + a)^n V$$

$$[1/F(D)] e^{ax} = [1/F(a)] e^{ax}$$

$$F(D^2) (a \sin mx + b \cos mx) = F(-m^2) (a \sin mx + b \cos mx)$$

$\int e^{ax} \cos(bx) \, dx$ and $\int e^{ax} \sin(bx) \, dx$ can be integrated by the 1/D method but it is simpler to consider the Real (or Complex) part of $\int e^{ax} [\cos(bx) + i \sin(bx)] \, dx = \int e^{(a+ib)x} \, dx = [1/(a + ib)] e^{(a+ib)x} + \text{constant}$

Integration by Parts

$d/dx (u v) = v \, du/dx + u \, dv/dx$, thus $\int u \, dv = uv - \int v \, du$ Use this formula to transform an Integral of a product

example (i) $\int x \sin(x) \, dx$ Put $x = u$ and $\sin(x) \, dx = dv$ hence $v = -\cos(x)$ and $du = dx$

example(ii) $\int x \ln(x) \, dx$ Put $\ln(x) = u$ and $x \, dx = dv$ hence $v = (\frac{1}{2}) x^2$ and $du = 1/x \, dx$

Standard forms

y	dy / dx	$\int y \, dx$
$a x^n$	$n a x^{n-1}$	$a x^{n+1} / (n + 1)$
a / x	$- a/x^2$	$a \ln(x)$
$\sin \omega x$	$\omega \cos \omega x$	$(-1/\omega) \cos \omega x$
$\cos \omega x$	$-\omega \sin \omega x$	$(1/\omega) \sin \omega x$
$\tan \omega x$	$\omega \sec^2 \omega x$	$-(1/\omega) \ln(\cos \omega x)$
$\sec x$	$\tan x \sec x$	$\ln(\sec x + \tan x)$
$\operatorname{cosec} x$	$-\cot x \operatorname{cosec} x$	$\ln(\operatorname{cosec} x - \cot x)$
$\cot x$	$-\operatorname{cosec}^2 x$	$\ln(\sin x)$
$\operatorname{Arc} \sin(x/a)$	$1 / \sqrt{a^2 - x^2}$	$x \operatorname{Arc} \sin(x/a) + \sqrt{a^2 - x^2}$
$\operatorname{Arc} \cos(x/a)$	$-1 / \sqrt{a^2 - x^2}$	$x \operatorname{Arc} \cos(x/a) - \sqrt{a^2 - x^2}$
$\operatorname{Arc} \tan(x/a)$	$a / (a^2 + x^2)$	$x \operatorname{Arc} \tan(x/a) - a \ln \sqrt{a^2 + x^2}$
e^{ax}	$a e^{ax}$	$(1/a) e^{ax}$
a^x	$a^x \ln(a)$	$a^x / [\ln(a)]$
$\ln(ax)$	$1/x$	$x [\ln(ax) - 1]$
$\log_a x$	$(1/x) \log_a e$	$x \log_a(x/e)$
$\sinh x$	$\cosh x$	$\cosh x$
$\cosh x$	$\sinh x$	$\sinh x$
$\tanh x$	$\operatorname{sech}^2 x$	$\ln(\cosh x)$
$\operatorname{Arc} \sinh(x/a)$	$1 / \sqrt{a^2 + x^2}$	$x \operatorname{Arc} \sinh(x/a) - \sqrt{a^2 + x^2}$
$\operatorname{Arc} \cosh(x/a)$	$1 / \sqrt{x^2 - a^2}$	$x \operatorname{Arc} \cosh(x/a) - \sqrt{x^2 - a^2}$
$\operatorname{Arc} \tanh(x/a)$	$a / (a^2 - x^2)$	$x \operatorname{Arc} \tanh(x/a) + a \ln \sqrt{a^2 - x^2}$

Functions of Time and other variables

Velocity $v = dx/dt$

Acceleration = $d^2x/dt^2 = v \, dv/dx$

Functions of two or more variables

$$V = F(x, y, z)$$

$$\delta V = (\partial V / \partial x) \delta x + (\partial V / \partial y) \delta y + (\partial V / \partial z) \delta z$$

where $\partial V / \partial x$ means the differential of V with respect to x while y and z are kept constant.


Areas and Volumes

Cones and Pyramids **Volume = (1/3) Base Area x Height**

Volume of Revolution,
ie volume enclosed by rotating a curve about the x axis

$$\text{Volume} = \int \pi y^2 dx$$

Spheres **Surface Area = $4 \pi R^2$** = Curved area of enclosing cylinder
Volume = $4/3 \pi R^3$



Maxima and Minima

$y = F(x)$ is a **Maximum** when $dy/dx = 0$ and d^2y/dx^2 is **negative**

$y = F(x)$ is a **Minimum** when $dy/dx = 0$ and d^2y/dx^2 is **positive**

$y = F(x)$ is a point of inflection when $dy/dx = 0$ and $d^2y/dx^2 = 0$

Graphs

Length of Arc $s = \int \sqrt{1 + (dy/dx)^2} dx$

Radius of Curvature $\rho = [1 + (dy/dx)^2]^{3/2} / (d^2y/dx^2)$

Vectors

Definitions;

Scalar has Magnitude but not Direction

Vector has magnitude and Direction

The Operator j rotates a vector 90° anticlockwise, $j^2 \mathbf{V} = -\mathbf{V}$, hence $j = \sqrt{-1} = i$, is one solution

The Operator h rotates a vector 120° anticlockwise, hence $h^3 \mathbf{V} = \mathbf{V}$ and $(1 + h + h^2) \mathbf{V} = 0$

Convention for axes. Shake Hands, Right Hand, Fingers point as i , Palm points as j , Thumb points as k

Matrix notation of a Vector. $| a_i \ a_j \ a_k |$ means Vector $a_i \mathbf{i} + a_j \mathbf{j} + a_k \mathbf{k}$

If $\mathbf{V} = V_x \mathbf{i} + V_y \mathbf{j} + V_z \mathbf{k}$ then $V = \sqrt{V_x^2 + V_y^2 + V_z^2}$

Let θ be the angle between two vectors \mathbf{V} and \mathbf{U}

$\mathbf{V} \cdot \mathbf{U} = V U \cos \theta$ hence $\cos \theta = [V_x U_x + V_y U_y + V_z U_z] / \sqrt{\{V_x^2 + V_y^2 + V_z^2\} \{U_x^2 + U_y^2 + U_z^2\}}$

\mathbf{V} and \mathbf{U} are orthogonal if $V_x U_x + V_y U_y + V_z U_z = 0$

$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$ where α, β and γ are the angles between a vector and each axis.

$\mathbf{V} \times \mathbf{U} = V U \sin \theta \mathbf{a}$ where \mathbf{a} is a unit vector orthogonal to \mathbf{V} and \mathbf{U}

If \mathbf{A}, \mathbf{B} and \mathbf{C} define three adjacent edges of a parallelepiped

then Volume = $\mathbf{A} \times \mathbf{B} \cdot \mathbf{C} = \mathbf{B} \times \mathbf{C} \cdot \mathbf{A} = \mathbf{C} \times \mathbf{A} \cdot \mathbf{B}$

$\mathbf{V} \cdot \mathbf{U}$ is a Scalar, $\mathbf{V} \times \mathbf{U}$ is a Vector, The differential of a Vector is a Vector

If a scalar value F is assigned to all points in a three dimensional volume, then Grad F (written ΔF) at any point is a Vector normal to the surface which connects the point to all points which have the same value of F .

ΔF has the magnitude equal to the differential of F with respect to distance in this direction

If a Vector \mathbf{F} is assigned to all points in a 3D volume,

Div \mathbf{F} is defined as $\partial \mathbf{F} / \partial x \cdot \mathbf{i} + \partial \mathbf{F} / \partial y \cdot \mathbf{j} + \partial \mathbf{F} / \partial z \cdot \mathbf{k} = \Delta \cdot \mathbf{F}$

and Curl \mathbf{F} is defined as $\partial \mathbf{F} / \partial x \times \mathbf{i} + \partial \mathbf{F} / \partial y \times \mathbf{j} + \partial \mathbf{F} / \partial z \times \mathbf{k} = \Delta \times \mathbf{F}$

Argand Diagram

The Complex Number $A + iB$ can be represented as a Vector $A + jB$


$A + iB = r [\cos \theta + i \sin \theta] = r e^{i\theta}$

where $r = \sqrt{A^2 + B^2}$ and $\theta = \text{Arc Tan}(B/A)$

Thus $[A + iB]^n = r^n e^{in\theta} = r^n [\cos(n\theta) + i \sin(n\theta)]$

Use for Multiplication or Division by Complex Numbers (and Vectors)

$\cos(2\pi n + \theta) + i \sin(2\pi n + \theta) = \cos \theta + i \sin \theta$ where n is any integer



Differential Equations

Definitions;

Ordinary or Partial (2 or more variables)

Order, if highest derivative is $d^n y / dx^n$ Order is n

Degree is the Index of highest derivative when rationalised

PI is the Particular Integral

CF is the Complementary Function

Complete Primitive = PI + CF

Singular Solution is an isolated solution

Linear Differential Equation has each term a Differential of y, all Degree one with Coefficient a function of x

(i) Solution of a Linear Differential Equation

$$\text{Put } y = a_0 + a_1 x + a_2 x^2/2! + a_3 x^3/3! + \dots + a_r x^r/r! + \dots$$

Check the answer has enough arbitrary constants

(ii) Exact Equations (first order) $Mdx + Ndy = 0$ This can be integrated immediately if $\partial N/\partial x = \partial M/\partial y$

(iii) Separate the variables to get $P(x) dx = Q(y) dy$ eg if $f(x) dy/dx = a$ then $y = \int [a/f(x)] dx + c$

(iv) Homogeneous Equations $dy/dx = f(y/x)$ Put $y = vx$

(v) Linear first order $dy/dx + P(x)y = Q(x)$ where P(x) and Q(x) are any function of x multiply by integrating factor $R = e^{\int P dx}$

(vi) Linear, constant coefficients $F(D)y = f(x)$ for example $7D^2(y) - 3D(y) + 9y = 2 + 3x$

CF Solve $F(D)y = 0$ by substituting $y = A e^{ax} + B e^{bx} + \dots$ where A, B, etc are arbitrary constants

Special cases (a) a and b conjugate pair $p \pm iq$, $y = e^{px} [A \cos(qx) + B \sin(qx)]$

(b) $a = b$, $y = A e^{ax} + B x e^{ax}$

PI Find one solution to $F(D) = f(x)$ and add to the CF to get the complete solution

Examples

(a) $f(x) = k_0 + k_1 x + k_2 x^2 + \dots$ Put $y = a_0 + a_1 x + a_2 x^2 + \dots$ Substitute and equate coefficients of x

(b) $f(x) = k \sin x$ or $k \cos x$ Put $y = a_1 \sin x + a_2 \cos x$ or take real (or complex) part of $y = A e^{ibx}$

(vii) Solution by Laplace Transform solves for f(t) and at the same time evaluates the arbitrary constants

This method is useful for evaluating the response of a control system

If $f(t) = A t^n e^{-at}$ then Laplace Transform $F(s) = A n! / (a + s)^{n+1}$

If $f(t) = A t^n \sin \omega t$ or $A t^n \cos \omega t$ then $F(s) = \text{Real or Complex part of } A n! / (s - j\omega)^{n+1}$

Laplace Transform of $d/dt [f(t)] = s F(s) - f(0)$

Laplace Transform of $d^2/dt^2 [f(t)] = s^2 F(s) - sf(0) - d/dt[f(0)]$

Laplace Transform of $\int f(t) dt = (1/s) F(s)$

(viii) $d^2 y/dx^2 = -A y$ This is SHM. Solve by multiplying by the integrating factor $2 dy/dx$

(ix) $x^2 d^2 y/dx^2 + x dy/dx + (x^2 - n^2) y = 0$

where $n = 0, 1, 2, 3, 4, \dots$ etc or $n = 1/2, 1/3, 1/4, \dots$ etc

This is Bessel's Eqn.

Solution is $y = A J_n(x) + B Y_n(x)$ where A and B are arbitrary constants

$J_n(x) = \sum [\{(-1)^s (x/2)^{2s+n} \} / \{ \Gamma(s+n+1) s! \}]$ from $s = 0$ to infinity

For +ive integers $\Gamma(x) = (x-1)!$ for other values, $\Gamma(x) = \int_0^{x-1} t^{x-1} e^{-t} dt$ from $t = 0$ to ∞

$Y_n(x) = [\cos n\pi J_n(x) - J_n(x)] / \sin n\pi$

$d^2 y/dx^2 + x y = 0$ can be converted to Bessel's Eqn by substitution

Fourier Series

Any cyclic function $y = F(x)$ can be converted to a series of the form

$$y = c_0 + a_1 \cos x + a_2 \cos 2x + \dots + a_n \cos nx + \dots \\ + b_1 \sin x + b_2 \sin 2x + \dots + b_n \sin nx + \dots$$

where

$$c_0 = \frac{1}{2\pi} \int_0^{2\pi} y dx$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} y \cos(nx) dx$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} y \sin(nx) dx$$

Summary (Part 2 Applied)

Mechanics

Constant acceleration equations

$$v = u + at \quad s = \left(\frac{1}{2}\right) (u + v) t \quad s = ut + \left(\frac{1}{2}\right) at^2 \quad v^2 = u^2 + 2as$$

Gravitational Force $F = G M_1 M_2 / d^2$

Moment of Inertia $I = \int x^2 dm$

Newton's Laws (summarised)

(i) A body moves in a straight line unless acted on by a force

(ii) $P = ma$ and $C = I d\omega/dt = I d^2\theta/dt^2$

(iii) Action and Reaction are equal and opposite

Conservation of Energy

$$\text{Work done} = F x = C \theta$$

$$\text{Kinetic Energy} = \left(\frac{1}{2}\right) m v^2 = \left(\frac{1}{2}\right) I \omega^2$$

$$\text{Potential Energy} = m g h$$

Conservation of Momentum

$$\text{Momentum} = m v \quad \text{Angular momentum} = I \omega$$

(u before, v after collision) $m_1 v_1 + m_2 v_2 = m_1 u_1 + m_2 u_2$
 $v_1 - v_2 = -e (u_1 - u_2)$

Body moving in a curve $\text{acceleration} = v^2/R = R \omega^2$

Two Dimension Forces in equilibrium $\text{Resultant Force in two directions} = 0$
 $\text{Plus Couple about any one point} = 0$

Three Forces in equilibrium are co-planar and either meet at a point or are parallel

Friction Force = μN under gravity $F = \mu mg$

Simple Harmonic Motion (SHM) $d^2x/dt^2 = -k x$

hence $x = a \sin \omega t + b \cos \omega t$ where $\omega = \sqrt{k}$

T the Time for one cycle (ie the Period) is given by $\omega T = 2\pi$, hence $\text{Period} = 2\pi / \sqrt{k}$

Capstan $P_2 = P_1 e^{\mu\theta}$

Structures $\text{Stress} = p/A$ $\text{Strain} = x / L$ $E = \text{Stress}/\text{Strain}$

Beam carrying a load $p / y = E / R = M / I$

where

p is stress at distance y from the Neutral Axis

E is Young's Modulus

R is radius of curvature

M is the bending moment


I is the 2nd moment of area about the Neutral Axis

Hoop stress = $pD / 2t$

Longitudinal stress = $pD / 4t$

where p is pressure, D is outside dia, t is wall thickness

Use the outside diameter to allow for radial compression in the shell



Cantilever Beam

Moment M at L

$$d = ML^2 / (2EI) \quad \theta = ML / (EI)$$

Load W at L

$$d = WL^3 / (3EI) \quad \theta = WL^2 / (2EI)$$


Distributed Load W

$$d = WL^3 / (8EI) \quad \theta = WL^2 / (6EI)$$

Suspension Bridge
Hanging chain

$$\text{Parabolic } y = w x^2 / (2 F)$$

$$\text{Catenary } y = c [\text{Cosh } (x/c) - 1]$$



Gyroscopes

$$\mathbf{C} = \boldsymbol{\omega} \times \mathbf{M}$$

Where \mathbf{C} is a couple expressed as a corkscrew vector

$\boldsymbol{\omega}$ is the angular velocity of precession expressed as a corkscrew vector

\mathbf{M} is the angular momentum of the flywheel expressed as a corkscrew vector